

New occurrence records and updated global distribution of *Limnias novemceras* (Rotifera: Flosculariaceae) using citizen-contributed and published data

Phuripong Meksuwan,¹ Mayte Martín,² Yurachat Meksuwan,³ Supiyani Maiphae^{4,5*}

¹Science Program (Biology), Faculty of Science and Technology, Phuket Rajabhat University, Phuket, Thailand; ²Independent Researcher, Barcelona, Spain; ³Faculty of Technology and Environment, Prince of Songkla University, Phuket Campus, Phuket, Thailand; ⁴Animal Systematics and Ecology Speciality Research Unit (ASESRU), Department of Zoology, Faculty of Science, Kasetsart University, Bangkok, Thailand; ⁵Biodiversity Center Kasetsart University (BDCKU), Kasetsart University, Bangkok, Thailand

Abstract

We report new occurrence records and an updated global distribution of *Limnias novemceras* Meksuwan, Jaturapruek & Maiphae, 2018 (Rotifera: Flosculariaceae) based on citizen-science observations and published data. Twenty-two records were compiled, 19 from social media, extending the species range from its presumed endemic area in southern Thailand to Nearctic, Neotropical, Afrotropical, Palearctic, Sino-Japanese, and Oriental realms. Morphological features were consistent across populations, while minor corona variation and ecological traits indicate behavioral flexibility.

Key words: Biogeography; biodiversity; zooplankton; Rotifera; citizen science.

Correspondence to: supiyani.m@ku.ac.th

Introduction

Knowledge in biodiversity, ecology, and the biogeography of organisms is typically generated through field research, in which a researcher or team visits the habitats of target species. These habitats may range from small ponds and forests to entire mountain ranges or extensive coastal areas, where organisms are studied and collected using a variety of sampling methods. Specimens are primarily identified based on their morphological characteristics, which are then compared with existing taxonomic descriptions to assign the appropriate scientific name and taxonomic classification. If the observed characters do not match any known species, researcher may propose a new species or even a new taxonomic category (Meiri and Mace, 2007; Braby *et al.*, 2024). In addition to taxonomic identification, researchers also document ecological and biological aspects such as microhabitat preferences, feeding behavior, mating patterns, and interspecific interactions (Jaturapruek *et al.*, 2018; García-Morales *et al.*, 2021). The scope and depth of these observations are guided by the specific research questions and objectives.

In recent decades, the era of digital technology, the Internet, and social media has transformed how biodiversity data are generated. While traditional field expeditions remain essential, these new tools have made the process more convenient and cost-effective by reducing the need for extensive travel to distant locations. Online platforms allow researchers and citizen sci-

tists to share photographs, videos, and ecological observations in real time, providing valuable records that complement field sampling. As a result, biodiversity research can now be conducted with lower budgets and wider geographic coverage than ever before (Atsumi *et al.*, 2024; Jansen *et al.*, 2024). Even so, as taxonomists, we emphasize that accurate identification of any group of organisms requires highly skilled specialists. In many cases, working on a single order, family, or even genus may demand a lifetime of dedicated study (Meksuwan *et al.*, 2015; Franch *et al.*, 2024; Jattupan *et al.*, 2024; Jaturapruek and Maiphae, 2025; Pokpongmongkol *et al.*, 2025). However, citizen observations are proving to be valuable, particularly for species with distinct and easily recognizable features, where the risk of misidentification is minimal.

Limnias novemceras is a sessile rotifer first described from a stream in Krabi Province, southern Thailand, where individuals were found attached to the roots of the aquatic plant *Hydrilla verticillata* (L.f.) Royle (Meksuwan *et al.*, 2018). Since its original discovery, no additional records were reported, leading to the assumption that the species was restricted to southern Thailand. This perception changed when multiple observations of the species began to appear on social media platforms. The objectives of the present study were therefore to compile photographs and videos from various online sources that clearly display the diagnostic features of *L. novemceras*, and to analyze these materials in order to reassess its geographic distribution, morphological variation, and substrate utilization.

Methods

Photographs and videos of *Limnias novemceras* were compiled from various social media platforms, including Facebook, Instagram, iNaturalist, Reddit, YouTube, and X (formerly known as Twitter). Other platforms were also searched but no records were found. Additionally, published research articles were reviewed for material identifiable as *L. novemceras*, including instances in which the specimens had been assigned to a different taxon.

Searches were conducted using both the search engines of each platform and general internet search engines. The keywords applied were “*Limnias*,” “*L. melicerta*,” and, in some cases, the broader term “rotifer”.

Specimens were identified as *L. novemceras* based on the diagnostic characters described by Meksuwan *et al.* (2018). Identification was confirmed when all three key traits were observed: i) the dorsal plate with nine distinct projections (Fig. 1C); ii) a dorsal gap of the corona wider than the tube opening (Fig. 1D); and iii) ventral antennae extending beyond the fully expanded corona (Fig. 1 A,C). When either the dorsal gap or the ventral antennae could not be clearly assessed, identification was classified as uncertain.

Results

A total of 33 social media posts and 3 research publications, including the original species description, were compiled in this study. Of these, Facebook (12) contributed the most posts, followed by Instagram (8), YouTube (7), X (4), iNaturalist (1), and Reddit (1). However, among the social media data, 11 posts were

traced to four original videos that had been reposted on different social media sites (LN9, LN10, LN11, and LN21). In addition, another 11 posts were interpreted as likely depicting the same populations that individual authors photographed or filmed within short intervals of time (LN2, LN3, LN16, and LN18). After accounting for these duplications, 19 independent records were retained for analysis of the global distribution of *L. novemceras* (Tab. 1). Of these, three were specifically labeled as *L. novemceras* by the content creator, and one by a commenter. In the following sections, we examine biological information on *L. novemceras* from these online sources, focusing on its biogeography, morphology, and ecological characteristics.

Biogeographical distribution

Based on the most recent information in the Rotifer World Catalog, *L. novemceras* had previously been reported only from its type locality (Jersabek *et al.*, 2025). Our findings, however, show that the species occurs on six zoogeographic realms, including the Nearctic, Neotropical, Afrotropical, Palearctic, Sino-Japanese, and Oriental realms (Holt *et al.*, 2013). These records span tropical, subtropical, and temperate regions, with no observations yet from Australia or Antarctica (Fig. 2).

Morphological variation

Specimens from different continents exhibited remarkably little morphological variation, indicating a high degree of consistency across populations. The three diagnostic traits -the structure of the dorsal plate, the wide dorsal gap of the corona, and the elongated ventral antenna- consistently matched the original description and illustrations. Most specimens recorded from social media appeared to be attached to filamentous algae (Fig. 1A).



Fig. 1. General features of *Limnias novemceras* under the microscope (*in vivo*). A) Specimen with a transparent, ringed tube (LN20). B) Specimen with an opaque tube covered with debris (LN20). C) Contracted specimen showing the dorsal plate with nine projections and long antennae (LN19). D) Specimen elevating its corona well beyond the tube opening (LN14). Specimens are not shown to scale.

Tab. 1. Summary of records of *Limnias novemceras*, including locations, social media sources, and additional information.

Code	Location	Country	Substrate	Source	Authors	Link
LN1	Small lagoon, San José	Uruguay	Detached from substrate	YouTube (2015)	@microuuguay	https://www.youtube.com/watch?v=i3Ww0je0guQ
LN2*	Not specified	Ukraine	Filamentous green algae and macrophyte	Facebook (2017) X (2017)	Sergey Lysak (Сергей Лысак)	https://www.facebook.com/groups/Amateur.Microscopy/permalink/807754042739346/ https://x.com/digiP2eam/status/936137439470551040/photo/1 https://x.com/digiP2eam/status/929742267488919552/photo/1 https://x.com/digiP2eam/status/92853825846353889/photo/2
LN3*	Not specified	Ukraine	Filamentous green algae and macrophyte	Facebook (2018)	Sergey Lysak (Сергей Лысак)	https://www.facebook.com/groups/Amateur.Microscopy/permalink/867959546718795/
LN4	Stream, Krabi Province	Thailand	Leaves of <i>Hydrilla verticillata</i> (L.f.) Royle	X (2018) Meksuwan <i>et al.</i> (2018)	Phuripong Meksuwan, Rapeejan Jaturapruek & Suptayanit Maiphae	https://x.com/digiP2eam/status/976355001915277312/photo/1 https://zokeys.pensoft.net/articles.php?id=28098
LN5*	Not specified	Taiwan	Unidentified macrophyte	Facebook (2018)	Wei-Fang Tsai	https://www.facebook.com/groups/324721808316035/permalink/345899052864977/?mibextid=rS40aB7S9Ucbxw6v
LN6*	Not specified	Taiwan	Filamentous green algae	Facebook (2018)	Hei Xiaoxin	https://www.facebook.com/ever.ritchang/videos/10157060320132764/?idorvanity=324721808316035&mibextid=rS40aB7S9Ucbxw6v
LN7	Cement pond, Phuket Province	Thailand	Roots of <i>Pistia stratiotes</i> L.	Meksuwan and Somkhaoyai (2019)	Phuripong Meksuwan & Aphitsara Somkhaoyai	https://drive.google.com/file/d/1giWPW5i0DdW1VkmYnwscEcnltnYwUn/view?usp=sharing
LN8*	Lake, Xochimilco	Mexico	Roots of <i>Eichhornia crassipes</i> (Mart.) Solms (water hyacinth)	Jiménez-Santos <i>et al.</i> (2019)	Marco Antonio Jiménez-Santos, S. S. S. Sarma, S. Nandini	https://www.jeb.co.in/index.php?page=abstract&issue=201907_jul19&number=19
LN9	Not specified	Taiwan	Filamentous green algae	YouTube (2019) Facebook (2019)	@changlynn8730	https://www.youtube.com/shorts/QPoXo2vjF2A https://www.facebook.com/groups/324721808316035/permalink/436361567152058/?mibextid=rS40aB7S9Ucbxw6v
LN10	Not specified	Mexico	Unclear	YouTube (2020) Facebook (2020) Instagram (2020)	Uriel Ruiz	https://www.youtube.com/watch?v=ams4X8oG78U https://www.facebook.com/uriel.ruiz.710/videos/1535092820023328/?idorvanity=254539018060854&mibextid=rS40aB7S9Ucbxw6v https://www.instagram.com/p/CJMeEL5JEUP/?hl=es-es
LN11	Lake, Baltimore	USA	Detached from unclear substrate	YouTube (2020) Facebook (2020)	Steve Cunningham	https://www.youtube.com/watch?v=2SMshBeYhSU
LN12	Not specified	Not specified	Unidentified macrophyte	YouTube (2021)	@Omicrosis	https://www.facebook.com/groups/Amateur.Microscopy/permalink/1420071244840953/
LN13	Not specified	Not specified	Filamentous green algae	YouTube (2022)	@Omicrosis	https://www.youtube.com/watch?v=ZrQlpKuwlw(at aout 1:22-1:35 in the clip)
LN14	Shrimp tank	Japan	Not seen	Instagram (2022)	@suziebi3	https://www.youtube.com/watch?v=arPvZyRW8k&ab_channel=Omicrosis
LN15*	Benoni, Gauteng Province	South Africa	Detached from substrate	iNaturalist (2022)	Jane Trembath	https://www.instagram.com/p/CeEz9kmbBcX/?hl=es-es
LN16	Aquarium	Malaysia	Decomposing macrophyte	Facebook (2022)	Chen Kai Wen	https://mexico.inaturalist.org/photos/199427842
LN17*	Not specified	Taiwan	Not seen	Facebook (2022)	Zhan Zhengwei	https://www.facebook.com/groups/Amateur.Microscopy/posis/1988729997975072/ https://www.facebook.com/100033718688968/videos/473600310973021/
LN18	Artificial lake, Veneto Region	Italy	Filamentous algae	Instagram (2023)	@itsallinmyhands_micro	https://www.facebook.com/groups/324721808316035/permalink/1315794522542087/?mibextid=rS40aB7S9Ucbxw6v
LN19	Artificial pond, Barcelona	Spain	Filamentous green algae	Instagram (2024)	Mayte Martín	https://www.instagram.com/p/CvZqPc_NMQk/?hl=es-es https://www.instagram.com/p/Cy31gEGCq3/?hl=es-es
LN20	Artificial pond, Barcelona	Spain	Filamentous green algae	Instagram (2024)	Mayte Martín	https://www.instagram.com/reel/C_2OLBTNxL5/?utm_source=ig_web_copy_link&igsh=MzRlODBiNWFlZA==
LN21*	Lake, Southern New Jersey	USA	Filamentous green algae	Reddit (2024) YouTube (2024)	microscopequestion	https://www.instagram.com/reel/DDjdpSqAmSP/?utm_source=ig_web_copy_link&igsh=MzRlODBiNWFlZA== https://www.reddit.com/r/microscopy/comments/1equbhh/found_some_limnias_rotifers/ https://youtu.be/Kk7grE0ewk
LN22*	Not specified	Not specified	Unidentified decomposing substrate	Facebook (2024)	Rylee Alland	https://www.facebook.com/groups/Amateur.Microscopy/posis/280142504038893/

*Records with uncertain identification.




Fig. 2. Distribution of *Limnias novemceras*. Records without specified locations (LN12, LN13, and LN22*) are not shown on the map. The asterisk indicates records with uncertain identification.

Some populations were also observed constructing opaque tubes covered with debris (Fig. 1B). Egg production was noted in several social media records, with most individuals carrying one or two eggs (Fig. 1D). Notably, six specimens (LN10, LN11, LN14, LN18, LN19, and LN20) were observed extending their corona well above the tube opening, with LN14 occurring at the highest elevation (Fig. 1D).

Discussion

The results expand the known range of *L. novemceras*, showing that it occurs across six zoogeographic realms. This broad geographic range indicates that *L. novemceras* is not a narrowly endemic rotifer but can persist in diverse freshwater ecosystems. Its presence across both hemispheres and multiple continents suggests high dispersal potential, possibly facilitated by passive vectors such as aquatic birds, water currents, or human-mediated transport (Fontaneto, 2019; Green *et al.*, 2023). Moreover, these observations imply that *L. novemceras* possesses broad ecological tolerance, which may explain its success in colonizing diverse habitats ranging from warmer low-latitude waters to cooler mid-latitude regions. The absence of records from Australia and

Antarctica may reflect genuine biogeographic limits or a lack of targeted surveys or published observations in these regions, consistent with the “rotiferologist effect”, wherein apparent patterns of species richness are strongly shaped by where specialists have worked rather than by true biodiversity (Fontaneto *et al.*, 2012).

In addition to its wide distribution, morphological traits were generally consistent with the original description, supporting their reliability as diagnostic characters. However, variation in the degree of corona elevation indicates behavioral flexibility. This variation may be influenced by factors such as activity level, the presence of suspended particles in the surrounding water, individual health condition, or may simply represent normal behavioral variation (Obertegger and Wallace, 2023). Importantly, this study highlights the value of citizen observations, which can substantially broaden knowledge of morphological variation and even contribute to taxonomic research (Chandler *et al.*, 2017).

Beyond morphology, *L. novemceras* also exhibits behaviors that have long been recognized in sessile rotifers. The construction of opaque tubes covered with debris, observed in some populations, is a behavior recognized in tube-building sessile rotifers for more than a century (Hudson and Gosse, 1889, as cited in Wallace *et al.*, 2006). The debris consists of suspended particles from the water column and/or fecal material expelled by the rotifers. These

particles could be gathered by the ciliary currents of the corona and bound with secretions produced from epidermal glands (Franch-Meneu and Pastor, 2025). In addition, the ultrastructure of these tubes has been examined in detail with transmission electron microscopy (TEM), both in earlier work by Yang and Hochberg (2018) and more recently by Franch-Meneu and Pastor (2025).

With regard to habitat, although most specimens recorded from social media appeared to be attached to filamentous algae, we hypothesize that this pattern does not necessarily reflect a true preference for algal substrates over submerged parts of aquatic plants, which are the more typical attachment sites of sessile rotifers (Meksuwan *et al.*, 2011, 2014). Instead, it may result from the collection methods used by observers, who often scoop water or employ plankton nets to obtain samples. Portions of this material are subsequently examined under a microscope, where individuals of *L. novemceras* are observed attached to suspended substrates such as algal filaments and decayed plant fragments. Such an approach likely underestimates the true density of sessile rotifers in their natural habitats. A more suitable sampling protocol involves collecting submerged portions of aquatic plants, placing them in containers filled with filtered water from the habitat, and transporting them to the laboratory. Small plant fragments can then be examined in chambers filled with field water under a stereoscopic microscope. Once the target species is located, fresh slides can be prepared for detailed examination under a compound microscope, particularly for species-level identification (Meksuwan *et al.*, 2011). Following this method allows for a more accurate assessment of both habitat preferences and population density.

Reproductive observations also provide insights into the species' biology. Most individuals carried a single egg within the tube and occasionally two, consistent with the original species description (Meksuwan *et al.*, 2018). However, one record mentioned that three eggs were observed, although this was not shown in the post (LN18). Moreover, because amictic and mictic females of *L. novemceras* may differ in their egg production (Wallace *et al.*, 2006), the reproductive biology of this species remains an important subject for future study.

Finally, valuable data were obtained not only from original posts but also from public comments, which are particularly important in citizen-science research. In the present study, ecologists and rotifer taxonomists contributed noteworthy input through comments, providing species identifications and ecological observations. Two-way communication between posters and public users facilitated question-and-answer discussion, yielding more detailed information on the topic of interest. In several cases, comments also provided details on sample locations, habitat characteristics, and permissions to use posted photographs and video clips.

Notes on the limitations of data from social media

A common limitation in species distribution studies using citizen-generated data is the potential for misidentification by non-experts. In the present study, this issue was minimized because all rotifer records were verified and confirmed by experts based on clearly defined diagnostic traits described in this paper. Images or videos lacking sufficient morphological details for species-level confirmation were conservatively categorized as “uncertain.”

Another limitation often raised concerns the temporal accuracy of observations. Because the sampling date is frequently unavailable in social media posts, the upload date was used as a proxy in this study. This approach is considered appropriate, as our aim was

to document the geographical occurrence of *L. novemceras* rather than to analyze temporal patterns.

A further limitation involves the accuracy of geographic information. Some social media records lacked explicit geolocation data and were therefore inferred from post descriptions or user profiles. For instance, records from the Facebook group “Taiwan Freshwater Microorganism Fans Club” were assumed to originate from Taiwan. Although the spatial precision of these data varied - from exact localities to broader regional assignments- such resolution is sufficient for illustrating the broad distribution of *L. novemceras* across continents.

Overall, despite these inherent limitations, the verified and curated citizen-sourced records provided a reliable foundation for assessing the species' global distribution.

Conclusions

This study provides new occurrence records and an updated global distribution of *Limnias novemceras*, extending its range from southern Thailand to five continents. Citizen-science observations, combined with published data, confirm the consistent morphology and reveal ecological traits linked to tube construction and substrate use. These records highlight the value of citizen-contributed material for documenting and expanding knowledge of microscopic animal distributions.

Acknowledgements

We would like to thank the anonymous reviewers for their valuable comments and suggestions. We also gratefully recognize the contributions of citizen scientists worldwide, whose photographs and videos shared on social media platforms made this study possible. This study was financially supported by the International SciKU Branding (ISB), Faculty of Science, Kasetsart University.

References

- Atsumi K, Nishida Y, Ushio M, Nishi H, Genroku T, Fujiki S, 2024. Boosting biodiversity monitoring using smartphone-driven, rapidly accumulating community-sourced data. *eLife* 13:RP93694.
- Braby MF, Hsu Y, Lamas G, 2024. How to describe a new species in zoology and avoid mistakes. *Zool J Linn Soc* 202:zlae043.
- Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, et al., 2017. Contribution of citizen science towards international biodiversity monitoring. *Biol. Conserv* 213:280-294.
- Fontaneto D, Barbosa AM, Segers H, Pautasso M, 2012. The ‘rotiferologist’ effect and other global correlates of species richness in monogonont rotifers. *Ecography* 35:174-182.
- Fontaneto D, 2019. Long-distance passive dispersal in microscopic aquatic animals. *Mov Ecol* 7:10.
- Franch V, Meksuwan P, Wallace RL, 2024. The dorsal plate is a critical feature in the reassessment of the rotiferan genus *Ptygura* (Monogononta; Gnesiotrocha; Flosculariidae). *Zootaxa* 5428:107-123.
- Franch-Meneu V, Pastor LM, 2025. Ultrastructure of the epidermis and tube of two sessile species of the genus *Limnias* (Rotifera:

- Monogononta: Gnesiotrocha: Flosculariidae). *Zoomorphology* 144:29.
- García-Morales AE, Domínguez-Domínguez O, Elías-Gutiérrez M, 2021. Uncovering hidden diversity: Three new species of the *Keratella* genus (Rotifera, Monogononta, Brachionidae) of high altitude water systems from central Mexico. *Diversity* 13:676.
- Green AJ, Lovas-Kiss Á, Reynolds C, Sebastián-González E, Silva GG, van Leeuwen CHA, Wilkinson DM, 2023. Dispersal of aquatic and terrestrial organisms by waterbirds: A review of current knowledge and future priorities. *Freshwater Biol* 68:173-190.
- Holt BG, Lessard J-P, Borregaard MK, Fritz SA, Araújo MB, Dimitrov D, et al., 2013. An update of Wallace's zoogeographic regions of the world. *Science* 339:74-78.
- Jansen M, Beukes M, Weiland C, Blumer M, Rudolfi M, Poerting J, et al., 2024. Engaging citizen scientists in biodiversity monitoring: Insights from the WildLIVE! Project. *Citiz Sci Theory Pract* 9:6.
- Jattupan S, Jaturapruek R, Sa-artrit P, Inuthai J, Ngernsaengsaruy C, Maiphae S, 2024. Diversity and habitat preferences of bdelloid rotifers in mosses and liverworts from beach forest along sand dunes in Thailand. *PeerJ* 12:e18721.
- Jaturapruek R, Maiphae S, 2025. An updated checklist of freshwater rotifers in Thailand: Taxonomy, diversity and distribution. *Biodivers Data J* 13:e155363.
- Jaturapruek R, Fontaneto D, Meksuwan P, Pholpunthin P, Maiphae S, 2018. Planktonic and periphytic bdelloid rotifers from Thailand reveal a species assemblage with a combination of cosmopolitan, tropical, and yet undescribed species. *Syst Biodivers* 16:128-141.
- Jiménez-Santos MA, Sarma SSS, Nandini S, 2019. Temperature-dependent demographic differences in sessile rotifers of the genus *Limnias* (Rotifera: Gnesiotrocha). *J Environ Biol* 40:711-718.
- Jersabek CD, Segers H, Leitner M, 2025. Rotifera World Catalog (RWC). *Limnias novemceras* Meksuwan, Jaturapruek & Maiphae, 2018. Accessed: 2025-09-03. Available from: <https://rotifera.aphia.org/aphia.php?p=taxdetails&id=1796390>.
- Meiri S, Mace GM, 2007. New taxonomy and the origin of species. *PLoS Biol* 5:e194.
- Meksuwan P, Pholpunthin P, Segers H, 2011. Diversity of sessile rotifers (Gnesiotrocha, Monogononta, Rotifera) in Thale Noi Lake, Thailand. *Zootaxa* 2997:1-18.
- Meksuwan P, Pholpunthin P, Walsh EJ, Segers H, Wallace RL, 2014. Nestedness in sessile and periphytic rotifer communities: A meta-analysis. *Int Rev Hydrobiol* 99:1-10.
- Meksuwan P, Pholpunthin P, Segers HH, 2015. Molecular phylogeny confirms Conochilidae as ingroup of Flosculariidae (Rotifera, Gnesiotrocha). *Zool. Scr.* 44(6):562-573.
- Meksuwan P, Jaturapruek R, Maiphae S, 2018. Two new species of genus *Limnias* (Rotifera, Gnesiotrocha) from Thailand, with keys to congeners. *Zookeys* 787:1-15.
- Obertegger U, Wallace RL, 2023. Trait-based research on Rotifera: The holy grail or just messy? *Wate.* 15:1459.
- Pokpongmongkol P, Jaturapruek R, Sa-artrit P, Maiphae S, 2025. Substrate preference and seasonal distribution of bdelloid rotifers in mosses in a primary forest in Thailand. *Diversity* 17:171.
- Wallace RL, Snell TW, Ricci C, Nogrady T, 2006. Rotifera, Volume 1: Biology, ecology and systematics (2nd ed.). Leide, Backhuys Publishers; 299 p.
- Yang H, Hochberg R, 2018. Ultrastructure of the extracorporeal tube and "cement glands" in the sessile rotifer *Limnias melicerta* (Rotifera: Gnesiotrocha). *Zoomorphology* 137:1-12.

Received: 31 October 2025; Accepted: 15 December 2025.

Edited by: Diego Fontaneto, *CNR-IRSA Water Research Institute, Verbania-Pallanza, Italy*.

Citation: Meksuwan P, Martín M, Meksuwan Y, Maiphae S. New occurrence records and updated global distribution of *Limnias novemceras* (Rotifera: Flosculariaceae) using citizen-contributed and published data. *J Limnol* 2026;85:2251.

Contributions: Phuripong Meksuwan, Supiyaniit Maiphae, conceptualization; Phuripong Meksuwan, Mayte Martín, Yurachat Meksuwan, data collection; Supiyaniit Maiphae, funding acquisition; Phuripong Meksuwan, Mayte Martín, Yurachat Meksuwan, Supiyaniit Maiphae, data analysis, writing – review and editing; Phuripong Meksuwan, Mayte Martín, Supiyaniit Maiphae, writing – original draft preparation. All authors read and approved the final manuscript.

Conflict of interest: the authors declare no competing interests, and all authors confirm accuracy.

Data availability: all data generated or analyzed during this study are included in this published article.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).